

# A Survey of Iranian Population Attitude About Vaccination Against COVID-19 and Evaluation of Its Spontaneous Reporting of Adverse Effects

Sevda Micaeili-Mirak<sup>1</sup>, Soha Namazi<sup>2,3</sup> and Elliyeh Ghadrdan<sup>4\*</sup>

<sup>1</sup>Department of Clinical Pharmacy, School of Pharmacy, Ardabil University of Medical Sciences, Iran

<sup>2</sup>Department of Clinical Pharmacy, Faculty of Pharmacy, Tehran University of Medical Sciences, Iran

<sup>3</sup>Research Center for Rational Use of Drugs, Tehran University of Medical Sciences, Iran

<sup>4</sup>Department of Clinical Pharmacy, School of Pharmacy, Alborz University of Medical Sciences, Iran

## ABSTRACT

The COVID-19 pandemic continues to claim victims around the world, and Iran is no exception from this ravage. Vaccine development led to growing optimism regarding control of COVID-19. Vaccine acceptance by the majority of the population is important for the success in controlling COVID-19. This study assessed the attitude of Iranian population toward COVID-19 vaccination. Also, the adverse effects of the COVID-19 vaccines on vaccinated individuals were reported. This cross-sectional study was conducted from May to July 2021 among Iranian population. The data was collected using the online researcher-made questionnaire. The questionnaire contained 16 questions in 3 sections.

The questionnaire was filled out by 916 people in total. The majority of participants had 40-65 years old (N:480, 52.4%). More than half of participants were from health care team (N:543, 59.3%). The majority of participants trusted the vaccine and said they would inject it (N:582, 63.5%). However more than 80% of participants preferred non-domestic vaccines over Iranian ones. Potential adverse effect following vaccination was the main concern of responders (N:576, 62.9%). The most common adverse effects reported by vaccine recipients were temporary fatigue and muscle pain (N:313, 71.95%). The level of education, academic area and area of practice in the health system were significantly associated with the positive response to COVID-19 vaccination (P-value 0.0001, 0.0001 and 0.004 respectively). According to the results of this study, Iranian health authorities should provide clear information about the safety and efficacy of COVID-19 vaccines, especially domestic types, to increase public confidence and awareness regarding vaccines.

**KEYWORDS:** COVID-19; Vaccination; Iranian population; Attitude; Adverse effect

## INTRODUCTION

Since the first cases of COVID-19, were identified more than a year ago. Approximately 220 countries were affected by the COVID-19 outbreak starting in December 2019. The effects of

COVID-19 vary from person to person. The majority of people who have COVID-19 infection will show mild to moderate symptoms and recover without hospitalization; however, some will undergo hospitalization or even die due to their illness [1]. More than

### Quick Response Code:



**Address for correspondence:** Elliyeh Ghadrdan, Assistant Professor, Department of Clinical Pharmacy, Faculty of Pharmacy, Alborz University of Medical Sciences, Iran

**Received:** October 08, 2021

**Published:** October 19, 2021

**How to cite this article:** Sevda Micaeili-M, Soha N, Elliyeh G. A Survey of Iranian Population Attitude About Vaccination Against COVID-19 and Evaluation of Its Spontaneous Reporting of Adverse Effects. 2021- 4(6) OAJBS.ID.000337. DOI: [10.38125/OAJBS.000337](https://doi.org/10.38125/OAJBS.000337)

190 million cases and 4.09 million deaths worldwide have been attributed to the COVID-19 pandemic as of July 17, 2021 [2]. Human health was not the only aspect of human life that was affected by this disease; economic conditions, and social relationships were also impacted [3]. It appears that the virus is transmitted mainly through the inhalation of droplets with a predominance of respiratory symptoms [4]. Several strategies were implemented to stop the spread of COVID-19, including social distancing, hand washing, face mask wearing, and reducing presence in crowded places [5]. Despite the fact that these methods have slowed disease spread, these methods are challenged by lack of public support or resilience, noncompliance with national and international guidelines, and inadequate infrastructure and resources [6].

As the disease had no specific treatment and many people died daily, the international community became focused on vaccination and increasing public immunity against it. Currently, a major public health measure that can save millions of lives worldwide is the vaccination. To date, different platforms including inactivated virus, recombinant protein, RNA based vaccine and deficient adenovirus based, have been used to manufacture vaccines [7]. As of 17 July 2021, human clinical trials have been conducted on 97 vaccines, and 32 vaccines have reached the final stages. Also, animal studies are ongoing on 77 preclinical vaccines [8]. As a result of the development of a vaccine against COVID-19, the human community gained hope for faster control of the disease. On the other hand, there are some concerns about safety, efficacy, and availability of COVID-19 vaccine among many people. Vaccine hesitancy remains a potential impediment to community uptake [9]. As defined by the WHO Strategic Advisory Group of Experts (SAGE), vaccine hesitancy is the refusal or delay of vaccination despite the availability of vaccination services [10]. The vaccine related adverse effects have a key role in public confidence in the vaccine. Fear of adverse effects is one of the reasons for people's distrust and rejection of the vaccine. The information on COVID-19 vaccines adverse effects comes from vaccine manufacturers' studies; so further observational studies

can provide more information on vaccines adverse effects. Raising knowledge about different aspects of COVID-19 vaccines can help allay concerns and improve public confidence [11].

With a population of about 84 million, the COVID-19 infection rate in Iran is high. As of 17 July 2021, more than 3.5 million COVID-19 confirmed cases and over 86966 deaths have been reported in Iran<sup>2</sup>. Several vaccines from different companies have entered the clinical studies phase. Currently in Iran two domestically made vaccines have been approved for emergency use from the Ministry of Health of Iran, there are also a number of vaccines in various stages of development in different domestic institutions [12]. It is important to examine the attitudes regarding COVID-19 vaccines among Iranian population as well as their acceptance of the vaccines. This study was conducted to evaluate the attitude and acceptance of Iranian population toward COVID-19 vaccination. A study of the adverse effects of the COVID-19 vaccine on vaccinated individuals was also conducted, in light of the fact that the occurrence of adverse effects is one of the concerns of people about the vaccination. However, several studies have examined the tendency to receive vaccines, preference for domestic versus non-Iranian vaccines [13], priority groups for vaccination [14], and adverse effects of vaccine [15] among special groups in Iran. These issues, along with other factors affecting attitudes, are comprehensively addressed in this study.

## MATERIALS AND METHODS

This survey is a cross-sectional study in which the attitude and acceptance of the Iranian population about COVID-19 vaccination were assessed. Vaccinated individuals were also evaluated for adverse effects of the vaccine. This study was conducted in August 2021. The study was reviewed and approved by the Ethics Committee of the Tehran University of medical sciences (IR.TUMS.TIPS.REC.1400.091). There was no consent form and completion of the questionnaire by each person was considered as her/ his satisfaction.

**Table 1:** The demographic characteristics of the study participants (N=916).

Variables		N	%
Gender	Male	237	25.9
	Female	679	74.1
Age (years)	<25	67	7.3
	25-39	348	38
	40-65	480	52.4
	>65	21	2.3
Level of education	Elementary	30	3.3
	Diploma or Associate degree	114	12.4
	Bachelor's degree or Master's degree	338	36.9
	Doctoral degree or post-doctoral degree	434	47.4
Academic area	Health-related	543	59.3
	Non-health related	373	40.7
Area of practice (If you are a member of the health care team)	Pharmacist	231	25.2
	Physician	99	10.8
	Dentist	78	8.5
	Nurse	48	5.2
	Paramedical sciences	49	5.3
	Other fields	38	4.1

**Table 2:** Attitude and acceptance toward COVID-19 vaccination.

Attitude and Acceptance	Items	N (%)
If it were possible to choose the type of COVID-19 vaccine, which type would you choose?	Iranian vaccine	158 (17.2)
	Non-domestic vaccine	758 (82.8)
Do you trust the COVID-19 vaccine, and do you inject it?	I trust/ I inject	582 (63.5)
	I trust/ I do not inject	16 (1.7)
	I do not trust/ I inject	217 (23.7)
	I do not trust/ I do not inject	88 (9.6)
	Due to catching COVID-19, I do not need a vaccine.	13 (1.4)
There is no need to use protective methods (e.g face mask use) after vaccination against COVID-19	Agree	73 (8)
	Disagree	751 (82)
	Neutral	92 (10)
What is the most important issue that concern you about COVID-19 vaccination?	Vaccine related adverse effects	576 (62.9)
	Anxiety about getting COVID-19 following vaccination	95 (10.4)
	Inadequate immunization following vaccination	344 (37.6)
	Lack of sufficient pharmaceutical data due to limited studies	293 (32)
	Confusion about which vaccine to choose among domestic and foreign manufacturers	249 (27.2)
	Injustice in the public distribution of vaccine	315 (34.4)
Which of the following groups is not a priority for COVID-19 vaccination?	Pregnant and lactating women	541 (59.1)
	Elderly people over 65 years of age	181 (19.8)
	Patients with chronic disease (e.g. cancer, diabetes and immunodeficiency)	79 (8.6)
	Healthcare providers	115 (12.6)
Would you recommend the COVID-19 vaccine to others?	Yes	791 (86.5)
	NO	123 (13.5)

**Table 3:** Evaluate adverse effects following COVID-19 vaccine administration.

Questions	Items	N (%)
Have you ever been vaccinated with the COVID-19 vaccine? Which one?	Yes/ Sputnik V	176 (19.2)
	Yes/ Sinopharm	23 (2.5)
	Yes/ AstraZeneca	200 (21.8)
	Yes/ Covaxin bhart biotech	15 (1.6)
	Yes/ Coviran barekat	0 (0)
	Yes/ Pfizer	6 (0.7)
	Yes/ Moderna	1 (0.1)
	Yes/ I do not know vaccine name	14 (1.5)
	No, I have not injected	481 (52.6)
After getting the vaccine, did you experience any adverse effects? Specify the type of complication.	Yes, I experienced adverse effect	327 (75.2)
	No, I did not have any problem after vaccine administration	108 (24.8)
How long did an adverse effect last following the injection of COVID-19 vaccine? N:435	Less than 24 hours	113 (25.9)
	24 - 72 hours	188 (43.2)
	3 - 7 days	22 (5.1)
	Over than 7 days	12 (2.8)
Did you take any action to improve this complication? N:435	No, I didn't take any special measures and the complication improved spontaneously	115 (26.4)
	Yes, I received chemical medicine like analgesics or antihistamines at home	222 (51.1)
	Yes, I received traditional medicine at home	2 (0.5)
	Yes, I was admitted to the hospital	1 (0.2)
If you experienced an adverse reaction after the first injection of the vaccine, do you administer the second dose?	Yes	345 (90.07)
	No	38 (9.93)

Data collection was conducted using the online researcher-made questionnaire. The questionnaire contained 16 questions in 3 sections (Table 1-3). The first part consisted of 5 questions about demographic information including age, sex, level of education, and occupational field. In 6 questions of second part, attitude and acceptance of general population and healthcare staffs regarding COVID-19 vaccine were evaluated. The third section of the study focused on those who had been vaccinated against COVID-19 and asked the participants about the occurrence of adverse effect after vaccination, its duration, and measures taken to eliminate it. This questionnaire was designed and then tested for validity and reliability, and then converted into an online form. The questionnaire was uploaded to online social media platforms (WhatsApp, Telegram and Instagram). There are more than 59 million internet users in Iran [16]. We sent a questionnaire to various social groups, and members of the groups were asked to respond to the questions and share the questionnaire with their contacts or acquaintances. In order to prevent duplicate answers, the questionnaire was designed to be answered only once by each participant. The content validity of the Persian-language questionnaire was evaluated by 9 experts (physicians, pharmacists and university professors). Content validity index and the content validity ratio calculated 0.96 and 0.82 respectively.

All data were analyzed using Statistical Package for the Social Sciences (SPSS) version 25. Shapiro-Wilk's test was utilized to assess the normality distributions of variables. Continuous variables were expressed as mean  $\pm$  standard deviation or median [interquartile range]. Independent sample t-test and Mann-Whitney U-test were performed to compare normal and non-

normal quantitative variables between the two groups. Categorical variables were presented as numbers and percentages. Chi-square and Fisher exact test were used to compare categorical variables. In order to assess the association of acceptance of the participants and other variables, the answers "I trust/ I inject" and "I do not trust/ I inject" were considered similar to yes and the options "I do not trust/ I do not inject" and "I trust/ I do not inject" and "Due to catching COVID-19, I do not need a vaccine" were considered similar to no and were tested by Chi-square/ Fisher exact test. In all comparisons, P-values less than 0.05 were considered as significant

The sample size was calculated with the Raosoft sample size calculator. In order to achieve a confidence level of 99% and margin of error of 5% for this study, a sample size of 664 people with a distribution of responses of 50% is required. A response rate of 30 to 40% is estimated for studies based on online questionnaires [17,18], so the questionnaire was uploaded in different social groups with a high number of members.

## RESULTS

A total of 1836 people visited this questionnaire, but 916 participants completed it. The majority of participants had 40-65 years old (N:480, 52.4%) and more than 74% (N:679) of them were female. Most participants had high level of academic education (N: 434, 47.4%) doctoral or post-doctoral degree). More than half of participants were from health care team (N:543, 59.3%), whereas pharmacists constituted the majority (N:231, 25.2%). The sociodemographic characteristics of the participants were obtained in Table1.

**Table 4:** COVID-19 vaccination acceptance and adverse effects among healthcare providers and general population in Iran.

Question	Item	Health care provider (543)		General population (373)		P- value
		N	%	N	%	
If it were possible to choose the type of COVID-19 vaccine, which type would you choose?	Iranian vaccine	68	12.5	90	24.1	0.0001
	Non-domestic vaccine	475	87.5	283	75.9	
Do you trust the COVID-19 vaccine, and do you inject it?	Due to catching COVID-19, I do not need a vaccine	5	0.9	8	2.1	0.0001
	I do not trust/ I do not inject	28	5.2	60	16.1	
	I do not trust/ I inject	126	23.2	91	24.4	
	I trust/ I inject	379	69.8	203	54.4	
	I trust/ I do not inject	5	0.9	11	2.9	
After getting the vaccine, did you experience any adverse effect?	I don't have any injection	144	26.5	337	90.3	0.0001
	Yes, I experienced adverse effect	307	56.5	20	5.4	
	No, I did not have any problem after vaccine administration	92	16.9	16	4.3	
There is no need to use protective methods (e.g face mask use) after vaccination against COVID-19	Agree	21	3.9	52	13.9	0.001
	Disagree	497	91.5	254	68.1	
	Neutral	25	4.6	67	18.0	
Which of the following groups is not a priority for COVID-19 vaccination?	Pregnant and lactating women	364	67.0	177	47.5	0.004
	Patients with chronic disease	32	5.9	47	12.6	
	Elderly people over 65 years of age	97	17.9	84	22.5	
	Doctors and nurses	50	9.2	65	17.4	

$P < 0.05$  is considered as significant.

In terms of acceptance of COVID-19 vaccines, among respondents, more than 80% (N: 758) preferred non-domestic vaccines over Iranian ones. Nevertheless, the majority of participants trusted the vaccine and said they would inject it (N:582, 63.5%). Most participants, whether general population (N: 294, 78%) or health care team (N:505, 93%), tend to inject vaccine; but the percentage of people who refuse to inject vaccine was significantly higher for the general population (N:79, 19%) than for medical staff (N:38, 6.1%). Besides, 86.5% (N: 791) reported they recommend COVID-19 vaccine administration to others. Potential adverse effects following vaccination (N:576, 62.9%), as well as the possibility of inadequate immunization (N:344, 37.6%), were the main concerns of participants regarding COVID-19 vaccination. Most participants were well-informed about COVID-19 vaccination; they knew group priorities for vaccination (N:541, 59.1%) and believed that vaccination would not eliminate the need for protective measures (e.g., face masks) (N:751, 82 %). Compared

to the general population, medical staff had more correct answers to questions dealing with vaccination priorities (67% vs 47.5%,  $P$ -value: 0.004) and need for protective measures after vaccination (91.5% vs 68.1%,  $P$ -value: 0.001). The comparison between health care providers and the general population is shown in Table 4.

In this study, almost half of participants (435 participants, 47.4%) received COVID-19 vaccine, AstraZeneca® vaccine was mostly given to them (N: 200, 21.8%). Approximately 75.2 percent of participants (N:327) experienced one or more adverse effects after receiving the vaccine. The most common adverse effects reported by vaccine recipients were fatigue and muscle pain (N:313, 71.95%). Most vaccine adverse effects usually resolved by analgesics or antihistamines (N: 222, 51.1%) within 24 to 72 hours (N: 188, 43.2%), and only 2.8% (N:12) of cases lasted longer than a week. Majority of respondents (more than 90%, N:345) who had received the first dose of the vaccine expressed a desire to receive the next dose.

**Table 5:** Participant's characteristics and association with COVID-19 vaccination acceptance.

Variable	Items	Yes (N%)	No (N%)	P
Gender	Female	586 (73.3)	93 (79.5)	0.094
	Male	213 (26.7)	24 (20.5)	
Age (years)	<25	52 (6.5)	15 (12.8)	0.1
	25-39	304 (38.0)	44 (37.6)	
	40-65	425 (53.2)	55 (47.0)	
	>65	18 (2.3)	3 (2.6)	
Level of education	Elementary	18 (2.3)	12 (10.3)	0.0001
	Diploma or Associate degree	94 (11.8)	20 (17.1)	
	Bachelor's degree or master's degree	285 (35.7)	53 (45.3)	
	Doctoral degree or post-doctoral degree	402 (50.3)	32 (27.4)	
Academic area	Health-related	505 (63.2)	38 (32.5)	0.0001
	Non-health related	294 (36.6)	79 (67.5)	
Area of practice (If you are a member of the health care team)	Physician	95 (18.8)	4 (10.5)	0.004
	Dentist	78 (15.4)	0 (0)	
	Pharmacist	214 (42.4)	17 (44.7)	
	Nurse	42 (8.3)	6 (15.8)	
	Paramedical field	43 (8.5)	6 (15.8)	
	Other fields	33 (6.5)	5 (13.2)	

$P < 0.05$  is considered as significant.

As shown in Table 5, level of education, academic area and area of practice in the health system were significantly associated with the positive response to COVID-19 vaccination ( $P$ -value 0.0001, 0.0001 and 0.004 respectively).

## DISCUSSION

This study was conducted to investigate the acceptance and attitude of Iranian population toward COVID-19 vaccines. Additionally, adverse reactions following the vaccination and how to control them were investigated in vaccinated individuals.

Many people around the world are concerned about the use of the COVID-19 vaccine because of the lack of global experience with it, doubts about its effectiveness and potential adverse effects. Among the 916 participants in this study, 87.2% were eager to

receive the vaccine. Surprisingly, 23.7% of respondents (N: 217) said that they still inject vaccines despite not trusting them. There have been a variety of studies conducted in different countries to determine the acceptability of the COVID-19 vaccine. In a global survey, the COVID-19 acceptance of 13426 people from 19 countries was assessed. The highest acceptance rates were found in China (88.62%), followed by Brazil (85.36%), and the lowest rates were found in the Russian population (54.85%) [19]. There were acceptance rates of 53.1% in Kuwait [20], 37.4% in Jordan [4], and 64.7% in Saudi Arabia [21] among Asian countries. The present study found that health care providers accept the COVID-19 vaccine more than the general population. It can be attributed to the higher level of knowledge in the field of COVID-19, possibility of further COVID-19 infection because of special occupational conditions among health care providers, as well as the direct observation of



morbidity and mortality of this disease. In the study of Fares et al, in line with our results, it was observed that factors such as direct contact with patients with COVID-19 and obtaining sufficient information about COVID-19 vaccine have a significant effect on increasing the acceptance of health care workers [22].

This study found that people may be concerned about vaccinations for a variety of reasons. Participants most commonly cited possible complications following vaccinations, poor vaccination efficacy, and the lack of enough scientific backing to make the vaccine. These results are in line with those reported by Pogue and colleagues. They evaluated the attitude of Americans toward their vaccination against COVID-19. According to their study, most people were concerned by the adverse effects of vaccines, uncertain efficacy, and the short duration of vaccine studies [23]. In El-Elimat 's study, vaccine safety concerns were cited as a significant reason for vaccine rejection in the Jordanian population [4]. The present study showed that a large percentage of participants were willing to receive the vaccine, but that most prefer the non-Iranian vaccine to the Iranian vaccine. Similarly, 73% of Iranian professors and researchers who participated in another study conducted in August 2020 reported their preference for non-Iranian vaccines [13]. The distrust may have been caused, in part, by the idea that domestic products are inferior to foreign counterparts, as vaccines did as well as many other domestic items. In addition, the misinformation

and worrying rumors about domestic vaccines sparked public concern. Increased transparency by vaccine manufacturers and government officials regarding the production and distribution of locally produced vaccines can lead to greater public confidence in the field of domestic COVID-19 vaccines.

There is a need to prioritize the population to receive the vaccine in Iran, in light of the large population and the lack of adequate access to vaccine for all members at the same time; those involved in the study were asked to select the priority groups. The results showed that patients with chronic disease, healthcare providers and elderly population were reported as prioritized groups for COVID-19 vaccination. These groups were currently prioritized for vaccination by Iranian health policymakers. The results were similar to those found by Moradi et al., who surveyed 878 Iranians about priority groups for vaccination [14]. This study asked the participants about wearing masks and attention to protective methods after receiving the vaccine. Most participants, especially healthcare providers, believed that it was necessary to wear a mask following vaccination against COVID-19. The center for disease control and prevention (CDC) now recommends that persons who have been vaccinated use masks to protect themselves, particularly against the new variant of COVID-19, and to avoid spread to others [24].




Figure 1: Incidence of COVID-19 vaccines related adverse effects.

In another part of the study, people who had received the COVID-19 vaccine were asked about the occurrence of adverse effects following vaccination and how to manage it. During the study period, more healthcare providers were vaccinated than other populations. Participants received the AstraZeneca® vaccine at a higher proportion than other vaccines. This was due to greater imports of AstraZeneca® vaccine than other brands at the time of this study. Approximately 75% of participants experienced adverse effects after receiving the COVID-19 vaccine, with fatigue and muscle pain, fever, and headache being the most common complaints, respectively. The reported adverse effects were similar to those reported by the CDC, which identified tiredness, headache, muscle pain and fever as common adverse effects [25]. Fortunately, participants did not report life-threatening complications or anaphylaxis. A few participants also reported uncommon but important symptoms such as psychotic symptoms (e.g., restlessness and mood fluctuations), angina, and arrhythmia although they were temporary (Figure 1). At this time, these complications are rare and only reported in some case reports [26,27]. Most participants reported improvement in adverse effects after taking analgesics or antipyretics for 24 to 72 hours. The number of cases that required hospitalization from vaccination adverse effects was very low in this study.

This study was performed before vaccine was administered to a high percentage of the Iranian population. Therefore, people's perceptions about the vaccine or the pattern of adverse effects coming from the vaccine may change as the vaccine is distributed further. Additionally, the study didn't include all groups of people because those without access to mobile phones, computers, and the internet were unable to participate.

## CONCLUSION

In conclusion, this study assessed the attitude of Iranian population toward COVID-19 vaccination. Also, the adverse effects of the COVID-19 vaccine on vaccinated individuals were evaluated. The majority of participants trusted the vaccine and said they would inject it. However more than 80% of participants preferred non-domestic vaccines. Although participants were worried about receiving the vaccine due to fears of negative adverse effects, fatigue and muscle pain were the most common adverse effects reported by participants. No life-threatening complications were reported as well. However, more studies need to be done with more participants in order to evaluate the adverse effects of COVID-19 vaccines.

## REFERENCES

- Nalbandian A, Sehgal K, Gupta A, Madhavan MV, McGroder C, et al. (2021) Post-acute COVID-19 syndrome. *Nat Med* 27(4): 601-615.
- (2021) Reported cases and deaths by country or territory.
- Haleem A, Javaid M, Vaishya R (2020) Effects of COVID-19 pandemic in daily life. *CMRP* 10(2): 78-79.
- El-Elimat T, AbuAlSamen MM, Almomani BA, Al-Sawalha NA, Alali FQ (2021) Acceptance and attitudes toward COVID-19 vaccines: A cross-sectional study from Jordan. *PLoS one* 16(4): e0250555.
- (2021) How to Protect Yourself & Others, Content source: National Center for Immunization and Respiratory Diseases.
- Maqbool A, Khan NZ (2020) Analyzing barriers for implementation of public health and social measures to prevent the transmission of COVID-19 disease using DEMATEL method. *Diabetes Metab Syndr* 14(5): 887-892.
- Nagy A, Alhatlani B (2021) An overview of current COVID-19 vaccine platforms. *Comput Struct Biotechnol J* 19: 2508-2517.
- Zimmer C, Corum J, Wee SL (2021) Coronavirus vaccine tracker US: The New York Times.
- Cordina M, Lauri MA, Lauri J (2021) Attitudes towards COVID-19 vaccination, vaccine hesitancy and intention to take the vaccine. *Pharmacy practice* 19(1): 2317-2317.
- Butler R, MacDonald NE (2015) Diagnosing the determinants of vaccine hesitancy in specific subgroups: The Guide to Tailoring Immunization Programmes (TIP). *Vaccine* 33(34): 4176-4179.
- Riad A, Pokorná A, Attia S, Klugarová J, Koščik M, et al. (2021) Prevalence of COVID-19 vaccine side effects among healthcare workers in the Czech Republic. *J Clin Med* 10(7): 1428.
- Islamic republic of Iran ministry of health and medical education.
- Abouee-Mehrzi A (2021) A national survey of Iranian academicians' attitudes towards covid-19 vaccination. *CJHR* 6(1): 21-28.
- Moradi N, Heydari S, Zarei Z, Arabloo J, Rezapour A, et al. (2021) Public views on priority groups for covid-19 vaccination: a survey from Iran. *SEMJ* 22(7): e113359.
- Babamahmoodi F, Saeedi M, Alizadeh-Navaei R, Hedayatzadeh-Omran A, Mousavi A, et al. (2021) Side effects and immunogenicity following administration of the sputnik v covid-19 vaccine among health care workers; an observational study in Iran. Preprint.
- Digital 2021: Iran.
- Fincham JE (2008) Response rates and responsiveness for surveys, standards and the journal. *Am J Pharm Educ* 72(2): 43-43.
- Regmi PR, Waithaka E, Paudyal A, Simkhada P, van Teijlingen E (2016) Guide to the design and application of online questionnaire surveys. *Nepal J Epidemiol* 6(4): 640-644.
- Lazarus JV, Ratzan SC, Palayew A, Gostin LO, Larson HJ, et al. (2021) A global survey of potential acceptance of a COVID-19 vaccine. *Nat med* 27(2): 225-228.
- Alqudeimat Y, Alenezi D, AlHajri B, Alfouzan H, Almokhaizeem Z, et al. (2021) Acceptance of a COVID-19 vaccine and its related determinants among the general adult population in Kuwait. *Med Princ Pract* 30(3): 262-271.
- Al-Mohaithef M, Padhi BK (2020) Determinants of COVID-19 vaccine acceptance in Saudi Arabia: a web-based national survey. *J Multidiscip Healthc* 13: 1657-1663.
- Fares S, Elmnyer MM, Mohamed SS, Elsayed R (2021) COVID-19 vaccination perception and attitude among healthcare workers in Egypt. *J Prim Care Community Health* 12.
- Pogue K, Jensen JL, Stancil CK, Ferguson DG, Hughes SJ, et al. (2020) Influences on attitudes regarding potential COVID-19 vaccination in the United States. *Vaccines* 8(4): 582.
- When You've Been Fully Vaccinated (2021) Center for disease control and prevention.
- Possible side effects after getting a COVID-19 vaccine (2021) Center for disease control and prevention.
- Jeet Kaur R, Dutta S, Charan J, Bhardwaj P, Tandon A, et al. (2021) Cardiovascular adverse events reported from COVID-19 vaccines: a study based on WHO database. *Int J Gen Med* 14: 3909-3927.
- Srinivasan KN, Sathyamurthy I, Neelagandan M (2021) Relation between COVID-19 vaccination and myocardial infarction -Casual or coincidental? *IHJ CVCR* 5(2): 71-74.